

Patient Information

Patient Name	Birth Date (Month DD, YYYY)	Body Weight
--------------	-----------------------------	-------------

Dosage Information

Dosage administration date (Month DD, YYYY)
Dosage (mg q 6 hr)

Specimen Collection Times

Infusion start time	:	<input type="checkbox"/> a.m. <input type="checkbox"/> p.m.
Infusion stop time/first specimen draw time (Typically two hours after infusion start)	:	<input type="checkbox"/> a.m. <input type="checkbox"/> p.m.
Second specimen draw time	:	<input type="checkbox"/> a.m. <input type="checkbox"/> p.m.
Third specimen draw time	:	<input type="checkbox"/> a.m. <input type="checkbox"/> p.m.
Fourth specimen draw time	:	<input type="checkbox"/> a.m. <input type="checkbox"/> p.m.

Four plasma specimens are required for this test

- The first specimen should be drawn immediately after termination of an intravenous infusion of 0.8 mg/kg busulfan. Additional specimens should also be drawn at 1 hour, 2 hours and 4 hours after termination of infusion.
- Draw blood in a green-top (sodium heparin) tube(s). Spin down and send 1.0 mL of sodium heparin plasma frozen in plastic vial.
- Label each specimen appropriately (exact time of draw).
- Complete all sections of form.
- Send all four specimens and form together under one order.

Questions? Phone **1-800-533-1710**